

## ABELIAN OR COMMUTATIVE GROUP

$(G, \times)$  is said to be abelian or commutative group if  $a \times b = b \times a, \forall a, b \in G$ . The group which are not abelian are called non-abelian or non-commutative.

## FINITE AND INFINITE GROUPS

group contains a finite number of elements, it is called a finite group. If the number of elements in a group is infinite, it is called an infinite group.

## ORDER OF A GROUP

The number of elements in a finite group is called the order of the group. It is denoted by  $O(G)$ .

An infinite group is called a group of infinite order.

## EXAMPLES OF GROUPS

- (i) The set  $\mathbb{Z}$  of integers is an infinite abelian group with respect to the operation of addition but  $\mathbb{Z}$  is not a group with respect to the multiplication.
- (ii) Let  $G = \{1\}$ , then  $G$  is an abelian group of order 1 with respect to multiplication.
- (iii) Let  $G = \{0\}$ , then  $G$  is an abelian group of order 1 with respect to addition.
- (iv) Let  $G = \{1, -1\}$ , then  $G$  is an abelian group of order 2 with respect to multiplication.

## Remarks

① When we say  $*$  is a binary operation defined on a non-empty set  $G$ , it implies that  $G$  is closed for the binary operation  $*$ , i.e.

$$a \in G, b \in G \Rightarrow a * b \in G \quad \forall a, b \in G$$

② A group is not simply a set, but it is an algebraic structure.

③ Because of the associativity, the parenthesis can be dropped in products of more than two elements of a group and instead of writing  $a * (b * c)$  or  $(a * b) * c$  we may simply write  $a * b * c$ . The associative law can be extended to any finite number of elements.

## Definition 2: - Group

A nonempty set of elements  $G$  is said to form a group if in  $G$  there is defined a binary operation, called the product and denoted by  $\cdot$ , such that

(i)  $a, b \in G$  implies that  $a \cdot b \in G$  (closed)

(ii)  $a, b, c \in G$  implies that  $a \cdot (b \cdot c) = (a \cdot b) \cdot c$   
(associative law)

(iii) There exists an element  $e \in G$  such that  $a \cdot e = e \cdot a = a$  for all  $a \in G$   
(the existence of an identity element in  $G$ )

(iv) For every  $a \in G$  there exists an element  $a^{-1} \in G$  such that  $aa^{-1} = a^{-1}a = e$   
(the existence of inverse in  $G$ )