


Benzoin condensation

The **benzoin addition** is an addition reaction involving two aldehydes. The reaction generally occurs between aromatic aldehydes or glyoxals.^{[1][2]} The reaction produces an acyloin. In the classic application benzaldehyde is converted to benzoin.^[3]

The benzoin condensation was first reported in 1832 by Justus von Liebig and Friedrich Wöhler during their research on bitter almond oil.^[4] The catalytic version of the reaction involving cyanide was developed by Nikolay Zinin in the late 1830s.^{[5][6]}


Contents

- [Reaction mechanism](#)
- [Scope](#)
- [See also](#)
- [References](#)
- [External links](#)



Reaction mechanism

The reaction is catalyzed by nucleophiles such as a cyanide or an N-heterocyclic carbene (usually thiazolium salts). The reaction mechanism was proposed in 1903 by A. J. Lapworth.^[7] In the first step in this reaction, the cyanide anion (as sodium cyanide) reacts with the aldehyde in a nucleophilic addition. Rearrangement of the intermediate results in polarity reversal of the carbonyl group, which then adds to the second carbonyl group in a second nucleophilic addition. Proton transfer and elimination of the cyanide ion affords benzoin as the product. This is a reversible reaction, which means that the distribution of products is determined by the relative thermodynamic stability of the products and starting material.




In this reaction, one aldehyde donates a proton and one aldehyde accepts a proton. Some aldehydes can only donate protons, such as 4-dimethylaminobenzaldehyde whereas benzaldehyde is both a proton acceptor and donor. In this way it is possible to synthesise mixed benzoins, i.e. products with different groups on each half of the product. However, care should be taken to match a proton donating aldehyde with a proton accepting aldehyde to avoid undesired homo-dimerization.

Scope

The reaction can be extended to aliphatic aldehydes with base catalysis in the presence of thiazolium salts; the reaction mechanism is essentially the same... These compounds are important in the synthesis of heterocyclic compounds. The analogous 1,4-addition of an aldehyde to an enone is called the Stetter reaction.

In biochemistry, the coenzyme thiamine is responsible for biosynthesis of acyloin-like compounds utilizing the benzoin addition. This coenzyme also contains a thiazolium moiety, which on deprotonation becomes a nucleophilic carbene.

The asymmetric version of this reaction has been performed by utilizing chiral thiazolium and triazolium salts. Triazolium salts were found to give greater enantiomeric excess than thiazolium salts.^[8] An example is shown below.^[9]



Scheme 2. An intramolecular benzoin addition

Since the products of the reaction are thermodynamically controlled, the **Retro Benzoin Addition** can be synthetically useful. If a benzoin or acyloin can be synthesized by another method, then they can be converted into the component ketones using cyanide or thiazolium catalysts. The reaction mechanism is the same as above, but it occurs in the reverse direction. This can allow the access of ketones otherwise difficult to produce.

See also

- [Aldol addition](#)
- [Acyloin condensation](#)
- [Stetter reaction](#)
- [Umpolung](#)

References

1. Menon, Rajeev S.; Biju, Akkattu T.; Nair, Vijay (2016). "Recent advances in N-heterocyclic carbene (NHC)-catalysed benzoin reactions" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4901930>). *Beilstein Journal of Organic Chemistry*. **12**: 444–461. doi:[10.3762/bjoc.12.47](https://doi.org/10.3762/bjoc.12.47) (<https://doi.org/10.3762%2Fbjoc.12.47>). PMC 4901930 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4901930>). PMID 27340440 (<https://pubmed.ncbi.nlm.nih.gov/27340440>).
2. Enders, Dieter; Niemeier, Oliver; Henseler, Alexander (2007). "Organocatalysis by N-Heterocyclic Carbenes". *Chemical Reviews*. **107** (12): 5606–5655. doi:[10.1021/cr068372z](https://doi.org/10.1021/cr068372z) (<https://doi.org/10.1021%2Fcr068372z>). PMID 17956132 (<https://pubmed.ncbi.nlm.nih.gov/17956132>).
3. Roger Adams, C. S. Marvel (1921). "Benzoin". *Organic Syntheses*. **1**: 33. doi:[10.15227/orgsyn.001.0033](https://doi.org/10.15227/orgsyn.001.0033) (<https://doi.org/10.15227%2Forgsyn.001.0033>).
4. Wöhler, Liebig; Liebig (1832). "Untersuchungen über das Radikal der Benzoesäure". *Annalen der Pharmacie*. **3** (3): 249–282. doi:[10.1002/jlac.18320030302](https://doi.org/10.1002/jlac.18320030302) (<https://doi.org/10.1002%2Fjlac.18320030302>). hdl:2027/hvd.hxdg3f (<https://hdl.handle.net/2027%2Fhvd.hxdg3f>).
5. N. Zinin (1839). "Beiträge zur Kenntniss einiger Verbindungen aus der Benzoylreihe" (<https://zenodo.org/record/1426941>). *Annalen der Pharmacie*. **31** (3): 329–332. doi:[10.1002/jlac.18390310312](https://doi.org/10.1002/jlac.18390310312) (<https://doi.org/10.1002%2Fjlac.18390310312>).
6. N. Zinin (1840). "Ueber einige Zersetzungprodukte des Bittermandelöls" (<https://zenodo.org/record/1426951>). *Annalen der Pharmacie*. **34** (2): 186–192. doi:[10.1002/jlac.18400340205](https://doi.org/10.1002/jlac.18400340205) (<https://doi.org/10.1002%2Fjlac.18400340205>).
7. Lapworth, A. (1904). "CXXII.—Reactions involving the addition of hydrogen cyanide to carbon compounds. Part II. Cyanohydrins regarded as complex acids" (<https://zenodo.org/record/2159947>). *Journal of the Chemical Society, Transactions*. **85**: 1206–1214. doi:[10.1039/CT9048501206](https://doi.org/10.1039/CT9048501206) (<https://doi.org/10.1039%2FCT9048501206>).

8. Knight, Roland; Leeper, F. (1998). "Comparison of chiral thiazolium and triazolium salts as asymmetric catalysts for the benzoin addition". *J. Chem. Soc., Perkin Trans. 1* (12): 1891–1894. doi:[10.1039/A803635G](https://doi.org/10.1039/A803635G) (<https://doi.org/10.1039%2FA803635G>).
9. D. Enders, O. Niemeier & T. Balensiefer (2006). "Asymmetric Intramolecular Crossed-Benzoin Reactions by N-Heterocyclic Carbene Catalysis". *Angewandte Chemie International Edition*. **45** (9): 1463–1467. doi:[10.1002/anie.200503885](https://doi.org/10.1002/anie.200503885) (<https://doi.org/10.1002%2Fanie.200503885>). PMID [16389609](#) (<https://pubmed.ncbi.nlm.nih.gov/16389609/>).

External links

- [Animation of the reaction mechanism](#) (https://commons.wikimedia.org/wiki/File:Benz_A_startAnimGif.gif)
-

Retrieved from "https://en.wikipedia.org/w/index.php?title=Benzoin_condensation&oldid=993668061"

This page was last edited on 11 December 2020, at 21:03 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.