



* Potential Energy diagram and Concept of activated complex :-

A/c to activated Complex theory (ACT), the bimolecular reaction between two molecules A₂ & B₂ progresses through the formation of the so-called activated complex which then decomposes to yield the product AB -



Threshold Energy :-

The minimum energy which the reactant molecules should possess so that they may react to produce the product is called threshold energy.


Activation energy :- (E_a)

Reactant molecules possess some energy initially also. The minimum extra energy given to the reactant molecules to make their energy equal to threshold energy is called Activation energy.


Thus,

$$\text{Activation energy} = \text{Threshold energy} - \text{Energy possessed by molecule.}$$

A Curve is obtained by potential energy vs reaction co-ordinate. Reaction coordinate represents the profile energy change when reactants change into products.




Exothermic reaction



Endothermic reaction.

* Rate data and Mechanism of a chemical Reaction :-

For thermodynamic formulation of the activated complex theory, let us consider a simple bimolecular reaction -



Where, $[AB^*]$ is called Activated Complex

k^* is the equilibrium constant between reactant and activated complex.

From classical mechanics, the energy of vibrations is given by - RT/N_A .

and from quantum mechanics, it is given by $h\nu$.

Where,

$$h\nu = RT/N_A$$

$$\nu = RT/N_A h$$

Thus, the reaction rate is given by—

$$-\frac{d[A]}{dt} = k k_2 [AB^*] = k (k_B T/h) [AB^*] \quad \text{--- ①}$$

where, the factor k , called transmission coefficient.

Now, at equilibrium—

$$K^* = \frac{[AB^*]}{[A][B]}$$

$$\text{or } [AB^*] = K^* [A][B] \quad \text{--- ②}$$

Now putting this value in eqⁿ ①

$$-\frac{d[A]}{dt} = (k_B T/h) K^* [A][B] \quad \text{--- ③}$$

Thus the rate constant k_2 may be expressed as—

$$k_2 = (k_B T/h) K^* \quad \text{--- ④}$$

The equilibrium constant K^* can be expressed in terms of $(\Delta G^\circ)^*$ called the standard Gibbs free energy of activation.

Thus,

$$(\Delta G^\circ)^* = -RT \ln K^*$$

$$\text{or } (\Delta G^\circ)^* = (\Delta H^\circ)^* - T(\Delta S^\circ)^*$$

Thus, we obtained

$$k^* = e^{-(\Delta G^\ddagger)/RT}$$

$$= e^{(\Delta S^\ddagger)/R} e^{-(\Delta H^\ddagger)/RT}$$

Hence, Eq^s - (4) becomes -

$$k_2 = \left(k_B T / h \right) e^{(\Delta S^\ddagger)/R} e^{-(\Delta H^\ddagger)/RT} \quad \text{--- (5)}$$

Eq^s - (5) is well known Eyring equation.

& $(\Delta S^\ddagger)^*$ = standard entropy of activation.

$(\Delta H^\ddagger)^*$ = standard enthalpy of activation.

From

 Dr. A. R. Gupta, (chemistry)

 U.S. College