

# Uniform Continuity :-

we suppose  $f$  is continuous for every value of  $x$  in  $[a, b]$ . It means that if  $x_0 \in [a, b]$  then given  $\epsilon > 0$ , there exists a positive number  $\delta$  such that  $|f(x) - f(x_0)| < \epsilon$  for every point  $x$  in the interval  $]x_0 - \delta, x_0 + \delta[$ . The number  $\delta$  will depend upon  $x_0$  as well as  $\epsilon$  and so we may write it symbolically as  $\delta(\epsilon, x_0)$ . Now suppose that we keep  $\epsilon$  fixed and vary  $x_0$ . Then for a given  $x_0$  there will correspond a value  $\delta$ . The set of values of  $\delta$  corresponding to values of  $x_0$  in  $[a, b]$ , may or may not have a non-zero lower bound. If this set of values of  $\delta$  has a non-zero lower bound, say  $\delta_0$ , then for every  $x_0$  in  $[a, b]$ , we have  $|f(x) - f(x_0)| < \epsilon$  for all  $x$  such that  $|x - x_0| < \delta_0$ . In such a case, we say that the function  $f$  is uniformly continuous in  $[a, b]$ .

(2)

Definition :- A function  $f$  is said to be uniformly continuous in  $[a, b]$  iff for a given arbitrary small positive number  $\epsilon$ , there can be found a number  $\delta$ , depending only on  $\epsilon$ , such that

$$x_1, x_2 \in [a, b] \text{ and } |x_1 - x_2| < \delta$$

$$\Rightarrow |f(x_1) - f(x_2)| < \epsilon$$

we have defined uniform continuity with reference to a finite closed interval  $[a, b]$ . But no such restriction is necessary for uniform continuity. The definition is essentially the same for intervals such as  $]a, b[$ ,  $]a, \infty[$

It should, however, be noted carefully that uniform continuity is a property associated with an interval and not with a single point.

It follows from the definition of uniform continuity that for a given  $\epsilon$ , the number  $\delta$  should be such that the condition

$$|f(x_1) - f(x_2)| < \epsilon$$

is satisfied for any two points  $x_1, x_2$  in  $[a, b]$  such that  $|x_1 - x_2| < \delta$ .

(3)

Theorem :- A function which is continuous in a closed and bounded interval  $[a, b]$  is uniformly continuous in  $[a, b]$ .

Proof:- By theorem [ If  $f$  is a continuous function on the closed interval  $[a, b]$ , then the interval can always be divided up into a finite number of sub-intervals such that given  $\epsilon > 0$

$$|f(x_1) - f(x_2)| < \epsilon$$

where  $x_1$  and  $x_2$  are any two points in the same sub-interval. ] the interval  $[a, b]$  can be divided up into sub-intervals

$$[a, x_1], [x_1, x_2], \dots, [x_{n-1}, b]$$

such that for any two points  $x, \beta$  in the same sub-interval, we have

$$|f(x) - f(\beta)| < \frac{\epsilon}{2} \quad \text{--- (1)}$$

let  $\delta$  be a positive number which does not exceed the least of the numbers

$$x_1 - a, x_2 - x_1, \dots, b - x_{n-1}.$$