

⑥ 24 Transitional epithelium is found in
 Thursday
 (i) Kidney (ii) Ureter
 (iii) Trachea (iv) Blood vessel
 Ans 6 - (ii)

⑦ Trachea is lined by -
 (i) ciliated epithelium
 (ii) Sensory epithelium
 (iii) Transitional epithelium
 (iv) Pseudostratified epithelium
 Ans 7 - (iv)

⑧ Stratified squamous epithelium is found -
 (i) Buccal cavity (ii) Oesophagus
 (iii) Stomach (iv) Intestine
 Ans 8 - (ii)

⑨ In our body ciliated epithelium is found in -
 (i) Trachea, neurocoel of C.N.S. & Oviduct
 (ii) Ureter & Vagina
 (iii) Bile duct & Small intestine.
 (iv) Eustachian tube & buccal cavity
 Ans 9 - (i)

20. The ciliated epithelium in our body may be found in -
 (i) Trachea (ii) Ureter
 (iii) Bile duct (iv) Intestine

MAY

Q1) Simple cuboidal epithelium are found in -

- (i) Buccal cavity
- (ii) Trachea
- (iii) Urinary tubules of kidney
- (iv) Ureters.

Ans 11 - (iii)

Q2) The germinal epithelium found in the ovary of mammals is -

- (i) Squamous epithelium
- (ii) cuboidal epithelium
- (iii) columnar epithelium
- (iv) Pseudo stratified epithelium

Ans 12 - (ii)

Q3) Ciliated epithelium in vertebrates is present -

- (i) Kidney & Stomach
- (ii) Buccal cavity & Oviduct
- (iii) Stomach & Urinary tubules
- (iv) Lymph vessels

Ans 13 - (ii)

Q4) The cellular layer of epidermis of skin consists of -

- (i) Glandular cell
- (ii) columnar cells
- (iii) A squamous stratified epithelium
- (iv) A complex stratified epithelium

Ans 14 - (iv)