


A *symmetry operation* is an action that leaves an object looking the same after it has been carried out. For example, if we take a molecule of water and rotate it by 180° about an axis passing through the central O atom (between the two H atoms) it will look the same as before. It will also look the same if we reflect it through either of two mirror planes, as shown in the figure below.



Each symmetry operation has a corresponding *symmetry element*, which is the axis, plane, line or point with respect to which the symmetry operation is carried out. The symmetry element consists of all the points that stay in the same place when the symmetry operation is performed. In a rotation, the line of points that stay in the same place constitute a *symmetry axis*; in a reflection the points that remain

unchanged make up a *plane of symmetry*.



The symmetry elements that a molecule may possess are:




1. E - the identity. The identity operation consists of doing nothing, and the corresponding symmetry element is the entire molecule. Every molecule has at least this element.
2. C_n - an n -fold axis of rotation. Rotation by $360^\circ/n$ leaves the molecule unchanged. The H_2O molecule above has a C_2 axis. Some molecules have more than one C_n axis, in which case the one with the highest value of n is called the *principal axis*. Note that by convention rotations are *counterclockwise* about the axis.
3. σ - a plane of symmetry. Reflection in the plane leaves the molecule looking the same. In a molecule that also has an axis of symmetry, a mirror plane that includes the axis is called a vertical mirror plane and is labeled σ_v , while one perpendicular to the axis is called a horizontal mirror plane and is labeled σ_h . A vertical mirror plane that

How can we help you?



bisects the angle between two C_2 axes called a dihedral mirror plane, σ_d .

4. i - a center of symmetry. Inversion through the center of symmetry leaves the molecule unchanged. Inversion consists of passing each point through the center of inversion and out to the same distance on the other side of the molecule. An example of a molecule with a center of inversion is shown below.



5. S_n - an n-fold improper rotation axis (also called a rotary-reflection axis). The rotary reflection operation consists of rotating through an angle $360^\circ/n$ about the axis, followed by reflecting in a plane perpendicular to the axis. Note that S_1 is the same as reflection and S_2 is the same as inversion. The molecule shown above has two S_2 axes.